


# Breast Cancer Survivors and Sexuality: A Review of the Literature Concerning Sexual Functioning, Assessment Tools, and Evidence-Based Interventions

Jacqueline Dow, BSN, RN, and Lisa Kennedy Sheldon, PhD, APRN, AOCNP®



© Steve Debenport/iStockphoto

**Background:** Most breast cancer survivors (BCSs) experience sexual dysfunction from treatment-induced hormonal changes, making sexual assessment an important component of survivorship care. However, because many oncology healthcare providers do not perform such assessment, a number of BCSs endure unidentified sexual dysfunction. Barriers to assessment faced by providers must be identified, and further research should examine the perspectives of BCSs and their partners regarding preferences for sexual assessment and interventions.

**Objectives:** The purpose of this literature review was to explore tools used to assess sexual functioning and evidence-based interventions used to treat sexual dysfunction in BCSs.

**Methods:** Articles published from 2003–2013 that reported on the effects of breast cancer and its treatment on sexual functioning in BCSs and located through online database searches (PubMed, the Cochrane Library, MEDLINE®, CINAHL®) were reviewed.

**Findings:** Sexual dysfunction is a common problem among BCSs. Using various assessment tools, providers should regularly assess sexual functioning in BCSs during and after treatment to improve quality of life for BCSs and their partners.

Jacqueline Dow, BSN, RN, is a medical cardiology unit floor RN at Brigham and Women's Hospital and Lisa Kennedy Sheldon, PhD, APRN, AOCNP®, is an assistant professor in the College of Nursing and Health Sciences at the University of Massachusetts, both in Boston. The authors take full responsibility for the content of the article. The authors did not receive honoraria for this work. The content of this article has been reviewed by independent peer reviewers to ensure that it is balanced, objective, and free from commercial bias. No financial relationships relevant to the content of this article have been disclosed by the authors, planners, independent peer reviewers, or editorial staff. Dow can be reached at jackiedow@gmail.com, with copy to editor at CJONEditor@ons.org. (Submitted May 2014. Revision submitted October 2014. Accepted for publication October 14, 2014.)

Key words: breast cancer survivor; sexual dysfunction; assessment

Digital Object Identifier: 10.1188/15.CJON.456-461

With more than 3.1 million women living in the United States with a history of invasive breast cancer, the community of breast cancer survivors (BCSs) is growing, making assessments that focus on the effects of treatment on patients extremely important (American Cancer Society, 2014). The National Comprehensive Cancer Network ([NCCN], 2015) recommends that male and female survivors be evaluated for sexual dysfunction. In BCSs, sexual dysfunction is common. A study by Brédart et al. (2011) determined that BCSs experienced sexual dysfunction significantly more often than did a comparative population of women without breast cancer, attributing it to psychological and treatment-related factors. In addition, a study by Alder et al. (2008) discovered that, on the Female Sexual Function Index (FSFI), which is a short tool used for assessing sexual functioning in women, about 68% of BCSs scored below the cutoff that differentiates sexual dysfunction

from normal function. In this study, BCSs had a higher rate of sexual dysfunction than healthy women, and among survivors with a history of chemotherapy treatment, about 90% also presented with a FSFI score below the cutoff, indicating poor sexual functioning (Alder et al., 2008). A 2010 online survey involving patients with breast cancer (N = 1,956) drawn from the membership of a national Australian breast cancer organization and conducted by Ussher, Perz, and Gilbert (2012) revealed that 78% (n = 1,427) of participants reported a decreased frequency of sexual encounters, and 64% (n = 1,151) noted a decline in sexual pleasure.

Findings from these studies and others demonstrate that sexual dysfunction is common among women with breast cancer. However, in the 2010 study by Ussher, Perz, and Gilbert (2013) involving patients with breast cancer, the authors found that only 41% of participants (n = 790) had received information about sexual functioning after diagnosis and treatment. In addition, 68%